

<b><u>Common Name</u></b>	<b><u>Botanical Name</u></b>	<b><u>Seed Germination Instructions</u></b>
Vine maple	<i>Acer circinatum</i>	30-60 days warm-moist stratification followed by 90-120 days cold-moist stratification. Sow outside in fall.
Common yarrow	<i>Achillea millefolium</i>	No pretreatment required. Sow in fall or spring on or just below the soil surface. Needs warm soil and light to germinate.
Vanillaleaf	<i>Achlys triphylla</i>	60 days cold-moist stratification. Sow outside in fall to late winter.
Columbia monkshood	<i>Aconitum columbianum</i>	60 days cold-moist stratification. Sow outside in fall to late winter.
Horse mint	<i>Agastache urticifolia</i>	30-60 days cold-moist stratification. Sow outside in fall to late winter.
Western joeiweed	<i>Ageratina occidentalis</i>	No pretreatment required. Sow outside in fall to early spring. Germination may be improved with 30 days cold stratification
California dandelion	<i>Agoseris grandiflora</i>	No pretreatment required. Sow outside in fall to early spring. Germination may be improved with 30 days cold stratification.
Tapertip onion	<i>Allium acuminatum</i>	30-60 days cold-moist stratification. Sow outside in fall to late winter.
Pacific mountain onion	<i>Allium validum</i>	60-90 days cold-moist stratification. Sow outside in fall to late winter.
Menzies' fiddleneck	<i>Amsinckia menziesii</i>	60 days cold-moist stratification. Sow outside in fall for best germination.
Sharptooth angelica	<i>Angelica arguta</i>	60 days cold-moist stratification. Sow outside in fall to late winter.
Woodland madia	<i>Ansiocarpus madioides</i>	30-60 days cold-moist stratification. Sow outside in fall to late winter.
Western columbine	<i>Aquilegia formosa</i>	60 days cold-moist stratification. Sow outside in fall to late winter.
California spikenard	<i>Aralia californica</i>	90-120 days cold-moist stratification. Sow outside in fall for best germination.
Goat's beard	<i>Aruncus dioicus</i>	30-60 days cold-moist stratification. Sow outside in fall to late winter. Plant on soil surface or lightly cover. Seed needs light to germinate.
Heartleaf milkweed	<i>Asclepias cordifolia</i>	60 days cold-moist stratification. Sow outside in fall to early spring.
Narrowleaf milkweed	<i>Asclepias fascicularis</i>	60 days cold-moist stratification. Sow outside in fall to early spring.
Showy milkweed	<i>Asclepias speciosa</i>	60 days cold-moist stratification. Sow outside in fall to early spring.
California harebell	<i>Asyneuma prenanthoides</i>	60 days cold-moist stratification. Sow in fall to late winter. Sow on soil surface or lightly cover. Seed needs light to germinate.
Deltoid balsamroot	<i>Balsamorhiza deltoidea</i>	90 days cold-moist stratification. Sow outside in fall.
Tall Oregon grape	<i>Berberis (Mahonia) aquifolium</i>	90 days cold-moist stratification. Sow outside in fall to late winter.
Cascade Oregon grape	<i>Berberis (Mahonia) nervosa</i>	60 days cold-moist stratification. Sow in fall to late winter.
California brome	<i>Bromus carinatus</i>	No pretreatment required. Sow outside in fall to early spring.
Chinook brome	<i>Bromus laevipes</i>	No pretreatment required. Sow outside in fall.
Tolmie's cats ear	<i>Calochortus tolmiei</i>	30-60 days cold-moist stratification. Sow in fall to late winter.
Marsh marigold	<i>Caltha leptosepala</i>	60-90 days cold-moist stratification. Sow in fall to late winter.
Large camas	<i>Camassia leichtlinii</i>	60-90 days cold-moist stratification. Sow in fall to late winter.
Deer brush	<i>Ceanothus integerrimus</i>	Soak in hot (not boiling) water for 24 hours, then 60-90 days cold-moist stratification. Sow outside in fall to late winter.
Birch leaf mountain mahogany	<i>Cercocarpus betuloides</i>	60 days cold-moist stratification. Sow outside in fall to late winter.
Soaproot	<i>Chlorogalum pomeridianum</i>	No pretreatment required. Sow outside in fall to early spring.
Fireweed	<i>Chamerion angustifolia</i>	60 days cold-moist stratification. Sow outside in fall to late winter.
Western redbud	<i>Cercis occidentalis</i>	Soak in hot (not boiling) water for 24 hours, then 60-90 days cold-moist stratification. Sow outside in fall to late winter.
Western clematis	<i>Clematis ligusticifolia</i>	30-60 days cold-moist stratification. Sow outside in fall to late winter.
Hound's tongue	<i>Cynoglossum grande</i>	30-60 days cold-moist stratification. Sow outside in fall to late winter.
Mountain larkspur	<i>Delphinium glaucum</i>	90-120 days cold-moist stratification. Sow outside in fall.
Meadow larkspur	<i>Delphinium nuttallianum</i>	90-120 days cold-moist stratification. Sow outside in fall.
Henderson's shooting star	<i>Dodecatheon hendersonii</i>	60-90 days cold-moist stratification. Sow outside in fall to late winter.
Squirreltail grass	<i>Elymus elymoides</i>	No pretreatment required. Sow outside in fall to early spring.
Blue wildrye	<i>Elymus glaucus</i>	No pretreatment required. Sow outside in fall to early spring.
Capitate sandwort	<i>Eremogone congesta</i>	60-90 days cold-moist stratification. Sow outside in fall to late winter.
Tall wooly buckwheat	<i>Eriogonum elatum</i>	60 days cold-moist stratification. Sow outside in fall to late winter.
Arrowleaf buckwheat	<i>Eriogonum compositum</i>	60 days cold-moist stratification. Sow outside in fall to late winter.
Barestem buckwheat	<i>Eriogonum nudum</i>	60 days cold-moist stratification. Sow outside in fall to late winter.
Sulphur flower buckwheat	<i>Eriogonum umbellatum</i>	60 days cold-moist stratification. Sow outside in fall to late winter.
Oregon sunshine	<i>Eriophyllum lanatum</i>	30 days cold-moist stratification. Sow outside in fall to late winter.

<b><u>Common Name</u></b>	<b><u>Botanical Name</u></b>	<b><u>Seed Germination Instructions</u></b>
Henderson's fawn lily	<i>Erythronium hendersonii</i>	90 days cold-moist stratification. Sow outside in fall.
Scarlet monkey flower	<i>Erythranthe (Mimulus) cardinalis</i>	No pretreatment required. Sow outside in fall to early spring. Seeds need light to germinate, surface sow.
Yellow seep monkey flower	<i>Erythranthe (Mimulus) guttata</i>	No pretreatment required. Sow outside in fall to early spring. Seeds need light to germinate, surface sow.
California poppy	<i>Eschscholzia californica</i>	30 days cold-moist stratification. Sow outside in fall or early spring.
Western goldentop	<i>Euthamia occidentalis</i>	No pretreatment required. Sow outside in fall to early spring.
California fescue	<i>Festuca californica</i>	30 days cold-moist stratification. Sow outside in fall or early spring.
Roemer's fescue	<i>Festuca roemeri</i>	30 days cold-moist stratification. Sow outside in fall or early spring.
Cascara	<i>Frangula purshiana</i>	60-90 days cold-moist stratification. Sow in fall to late winter.
Bluehead gilia	<i>Gilia capitata</i>	30 days cold-moist stratification. Sow outside in fall.
White rushlily	<i>Hastingsia alba</i>	60-90 days cold-moist stratification. Sow outside in fall to late winter.
Bigelow's sneezweed	<i>Helenium bigelovii</i>	30 days cold-moist stratification. Sow outside in fall to late winter.
Cowparsnip	<i>Heracleum maximum</i>	120 days cold-moist stratification. Sow outside in fall.
Alum root	<i>Heuchera micrantha</i>	60 days cold-moist stratification. Sow outside in fall. Seeds need light to germinate, surface sow.
Oceanspray	<i>Holodiscus discolor</i>	90-120 days cold-moist stratification. Sow outside in fall.
Broad leaved lotus	<i>Hosackia crassifolia</i>	Soak in hot (not boiling) water for 24 hours, then 60-90 days cold-moist stratification. Sow in fall to late winter.
Scarlet gilia	<i>Ipomopsis aggregata</i>	30 days cold-moist stratification. Sow outside in fall to early spring.
Red beardtongue	<i>Keckiella corymbosa</i>	30 days cold-moist stratification. Sow outside in fall or early spring.
Bush beardtongue	<i>Keckiella lemmonii</i>	30 days cold-moist stratification. Sow outside in fall or early spring.
Bolander's tarweed	<i>Kyhosia bolanderi</i>	30-60 days cold-moist stratification. Sow outside in fall to late winter.
Siskiyou lewisia	<i>Lewisia cotyledon</i>	30-60 days cold-moist stratification. Sow outside in fall to late winter.
Blue wild flax	<i>Linum lewisii</i>	30 days cold-moist stratification. Sow outside in fall or early spring.
California lomatium	<i>Lomatium californicum</i>	60 days cold-moist stratification. Sow outside in fall to late winter.
Fernleaf biscuitroot	<i>Lomatium dissectum</i>	60 days cold-moist stratification. Sow outside in fall.
Bigseed biscuitroot	<i>Lomatium macrocarpum</i>	60 days cold-moist stratification. Sow outside in fall to late winter.
Nineleaf biscuitroot	<i>Lomatium triternatum</i>	60 days cold-moist stratification. Sow outside in fall to late winter.
Common lomatium	<i>Lomatium utriculatum</i>	60 days cold-moist stratification. Sow outside in fall to late winter.
Pink honeysuckle	<i>Lonicera hispidula</i>	60-90 days cold-moist stratification. Sow outside in fall.
Sickelkeel lupine	<i>Lupinus albicaulis</i>	Soak in hot (not boiling) water for 24 hours, then 30 days cold-moist stratification. Sow outside in fall to early spring.
Silver bush lupine	<i>Lupinus albifrons</i>	No pretreatment required. Sow outside in fall to early spring.
Velvet lupine	<i>Lupinus leucophyllus</i>	Soak in hot (not boiling) water for 24 hours, then 30 days cold-moist stratification. Sow outside in fall to early spring.
Elegant tarweed	<i>Madia elegans</i>	No pretreatment required. Sow outside in fall to early spring.
Large false Soloman's seal	<i>Maianthemum racemosum</i>	Seed takes two years to germinate. Sow outside in fall. 6 months cold-moist stratification, followed by 4 months warm-moist stratification, followed by another 5 months cold-moist stratification.
Giant blazingstar	<i>Mentzelia laevicaulis</i>	60-90 days cold-moist stratification. Sow outside in fall.
Coyote mint	<i>Monardella odoratissima</i>	30 days cold-moist stratification. Sow outside in fall to late winter.
Coyote tobacco	<i>Nicotiana attenuata</i>	No pretreatment required. Sow outside, or start in a greenhouse in spring.
Indian tobacco	<i>Nicotiana quadrivalvis</i>	No pretreatment required. Sow outside, or start in a greenhouse in spring.
False turtlehead	<i>Nothochelone nemorosa</i>	60-90 days cold-moist stratification. Sow outside in fall.
Douglas' grasswidow	<i>Olsynium douglasii</i>	90-120 days cold-moist stratification. Sow outside in fall.
Siskiyou Mountains owl's clover	<i>Orthocarpus cuspidatus</i>	90 days cold-moist stratification. Sow outside in fall.
Indian warrior	<i>Pedicularis densiflora</i>	30 days cold-moist stratification. Sow in fall. Hemiparasite, needs host plant to thrive. Parasitic on members of the heath family, such as manzanita or madrone.
Siskiyou penstemon	<i>Penstemon anguineus</i>	60-90 days cold-moist stratification. Sow outside in fall to late winter.
Hot rock penstemon	<i>Penstemon deustus</i>	60-90 days cold-moist stratification. Sow outside in fall to late winter.
Mountain blue penstemon	<i>Penstemon laetus</i>	60-90 days cold-moist stratification. Sow outside in fall to late winter.
Small flowered penstemon	<i>Penstemon procerus</i>	60-90 days cold-moist stratification. Sow outside in fall to late winter.

<b><u>Common Name</u></b>	<b><u>Botanical Name</u></b>	<b><u>Seed Germination Instructions</u></b>
Western sweet coltsfoot	<i>Petasites frigidus</i>	No pretreatment required. Sow outside in fall to early spring.
Varileaf phacelia	<i>Phacelia heterophylla</i>	30 days cold-moist stratification. Sow outside in fall or early spring. Seeds need light to germinate, surface sow.
Tall phacelia	<i>Phacelia procera</i>	30 days cold-moist stratification. Sow outside in fall. Seeds need light to germinate, surface sow.
Mock orange	<i>Philadelphus lewisii</i>	90-120 days cold-moist stratification. Sow outside in fall.
Pacific ninebark	<i>Physocarpus capitatus</i>	30 days cold-moist stratification. Sow outside in fall or early spring.
Hooker's fairybells	<i>Prosartes hookeri</i>	60-90 days cold-moist stratification. Sow outside in fall.
Chokecherry	<i>Prunus virginiana</i>	90-120 days cold-moist stratification. Sow outside in fall.
Bitter cherry	<i>Prunus emarginata</i>	90-120 days cold-moist stratification. Sow outside in fall.
Golden currant	<i>Ribes aureum</i>	90 days cold-moist stratification. Sow outside in fall.
Wax currant	<i>Ribes cereum</i>	90-120 days cold-moist stratification. Sow outside in fall.
Sierra gooseberry	<i>Ribes roezlii</i>	90-120 days cold-moist stratification. Sow outside in fall.
Sticky currant	<i>Ribes viscosissimum</i>	90 days cold-moist stratification. Sow outside in fall.
Thimbleberry	<i>Rubus parviflorus</i>	30 days cold-moist stratification. Sow outside in fall or early spring.
Waxy coneflower	<i>Rudbeckia glaucescens</i>	30 days cold-moist stratification. Sow outside in fall or early spring.
Western coneflower	<i>Rudbeckia occidentalis</i>	30 days cold-moist stratification. Sow outside in fall or early spring.
Blue elderberry	<i>Sambucus nigra</i> spp. <i>caerulea</i>	60-90 days cold-moist stratification. Sow outside in fall.
Pacific blacksnakeroot	<i>Sanicula crassicaulis</i>	30-60 days cold-moist stratification. Sow outside in fall or early spring.
Lambstongue ragwort	<i>Senecio integerrimus</i>	30-60 days cold-moist stratification. Sow outside in fall to late winter.
Arrowleaf ragwort	<i>Senecio triangularis</i>	30-60 days cold-moist stratification. Sow outside in fall or early spring.
Oregon checkermallow	<i>Sidalcea oregana</i>	60-90 days cold-moist stratification. Sow outside in fall.
Bell catchfly	<i>Silene campanulata</i>	30-60 days cold-moist stratification. Sow outside in fall to late winter.
West coast Canada goldenrod	<i>Solidago elongata</i>	No pretreatment required. Sow outside in fall to early spring.
Western mountain ash	<i>Sorbus scopulina</i>	90-120 days cold-moist stratification. Sow outside in fall.
Douglas' spiraea	<i>Spiraea douglasii</i>	30-60 days cold-moist stratification. Sow outside in fall to late winter.
Subalpine spiraea	<i>Spiraea splendens</i>	30-60 days cold-moist stratification. Sow outside in fall to late winter.
Lemmon's needlegrass	<i>Stipa lemmonii</i>	90 days cold-moist stratification. Sow outside in fall.
Claspleaf twistedstalk	<i>Streptopus amplexifolius</i>	Seed takes two years to germinate. Sow outside in fall. 6 months cold-moist stratification, followed by 4 months warm-moist stratification, followed by another 5 months cold-moist stratification.
Leafybract aster	<i>Symphyotrichum foliaceum</i>	No pretreatment required. Sow outside in fall to early spring.
Fringe cups	<i>Tellima grandiflora</i>	30 days cold-moist stratification. Sow outside in fall or early spring.
Western false asphodel	<i>Triantha occidentalis</i>	60-90 days cold-moist stratification. Sow outside in fall.
Vinegarweed	<i>Trichostema lanceolatum</i>	60-90 days cold-moist stratification. Sow in fall to late winter.
California false hellebore	<i>Veratrum californicum</i>	30-60 days warm-moist stratification followed by 90-120 days cold-moist stratification. Sow outside in early fall.
Western verbena	<i>Verbena lasiostachys</i>	30-60 days cold-moist stratification. Sow outside in fall or early spring.
American vetch	<i>Vicia americana</i>	No pretreatment required. Sow outside in fall to early spring.
Narrowleaf mule's ears	<i>Wyethia angustifolia</i>	90 days cold-moist stratification. Sow outside in fall.
Beargrass	<i>Xerophyllum tenax</i>	90-120 days cold-moist stratification. Sow outside in fall.